

IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet NO₃_VOC41

Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hardcopy without explicit written permission. The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, (<http://iupac.pole-ether.fr>).

This data sheet last evaluated: June 2013; last change in preferred values: June 2013.



Rate coefficient data

$k/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	Temp./K	Reference	Technique/ Comments
<i>Absolute Rate Coefficients</i>			
$(2.4 \pm 0.7) \times 10^{-11}$	298	Martínez et al., 1999	DF-LIF (a)
$(3.6 \pm 0.7) \times 10^{-11}$	433		
<i>Relative Rate Coefficients</i>			
$(2.94 \pm 0.05) \times 10^{-11}$	294	Atkinson et al., 1985	RR (b)

γ -terpinene is 1-isopropyl-4-methyl-cyclohexa-1,4-diene.

Comments

- (a) NO₃ radicals ($6\text{--}30 \times 10^{11} \text{ molecule cm}^{-3}$) generated from reaction of F atoms (made in a microwave discharge through F₂/He) with HNO₃. Flow tube was operated at ~1.33 mbar (1 Torr) He. γ -terpinene was present at similar concentrations (1-3 fold) to NO₃. So that absolute NO₃ concentrations (derived by titration with tetramethylethane) were necessary to derive the rate coefficient.
- (b) 4000 L Teflon chamber at 980 mbar (735 Torr) of air. NO₃ was generated by the thermal decomposition of N₂O₅. γ -terpinene and 2-methyl-2-butene (reference reactant) were monitored by GC. Correction made to the γ -terpinene loss rate due to reaction with NO₂ was <1 %. The rate constant ratio, $k(\text{NO}_3 + \gamma\text{-terpinene}) / k(\text{NO}_3 + 2\text{-methyl-2-butene}) = 3.14 \pm 0.05$ is placed on an absolute basis by $k(\text{NO}_3 + 2\text{-methyl-2-butene}) = 9.37 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at 298 K (Atkinson and Arey, 2003).

Preferred Values

Parameter	Value	T/K
$k/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	2.9×10^{-11}	298
<i>Reliability</i>		
$\Delta \log k$	± 0.12	298

Comments on Preferred Values

The preferred value of the room temperature rate coefficient is based on the relative rate study of Atkinson et al. (1985) in which accurate determination of the reactant concentrations was not required. The error limits are expanded to reflect the necessity to correct for removal of γ -terpinene by reaction with NO_2 and the $\approx 20\%$ difference to the absolute rate measurement at the same temperature (Martínez et al., 1999). The difference between the rate coefficients obtained at 298 and 433 K (factor 1.5) was not considered large enough to warrant a more detailed investigation of the temperature dependence (Martínez et al., 1999).


There are no product studies of this reaction, though the large rate constant indicates that the reaction proceeds mainly via addition of NO_3 across a double bond to form a chemically activated nitrooxyalkyl radical. At pressures found in the troposphere this adduct will undergo collisional stabilization prior to reaction with O_2 to form a nitrooxyalkyl peroxy radical or decompose to release NO_2 . At atmospheric pressure the formation of the peroxy radical will generally dominate.

References

Atkinson, R., Aschmann, S. M., Winer, A. M., and Pitts, J. N., *Env. Sci. Tech.*, 19, 159-163, 1985.

Atkinson, R., and Arey, J., *Chem. Rev.*, 103, 4605-4638, 2003.

Martínez, E., Cabañas, B., Aranda, A., Martín, P., and Salgado, S., *J. Atmos. Chem.*, 33, 265-282, 1999.



Rate coefficients for $\text{NO}_3 + \gamma$ -terpinene