


# IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet NOx6

Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hardcopy without explicit written permission. The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, <http://iupac.pole-ether.fr>.

This data sheet last evaluated: 4<sup>th</sup> June 2009; last change in preferred values 4<sup>th</sup> June 2009.



$$\Delta H^\circ = -189.7 \text{ kJ mol}^{-1}$$

## Rate coefficient data

$k / \text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	Temp./K	Reference	Technique/ Comments
<i>Absolute Rate Coefficients</i>			
$2.0 \times 10^{-11} \exp[(107 \pm 8)/T]$	104-354	Streit et al., 1976	PLP (a)
$(2.8 \pm 0.6) \times 10^{-11}$	300		
$(2.4 \pm 0.1) \times 10^{-11}$	295	Amimoto et al., 1979	PLP-RA (b)
$(2.52 \pm 0.25) \times 10^{-11}$	297	Wine and Ravishankara, 1981	PLP-RF (c)
$2.2 \times 10^{-11} \exp[(118 \pm 21)/T]$	195-673	Blitz et al., 2004	PLP-LIF (d)
$(3.06 \pm 0.25) \times 10^{-11}$	295		
$1.99 \times 10^{-11} \exp[(145 \pm 9)/T]$	197-427	Strekowski and Wine, 2004	PLP-RF (e)
$3.24 \times 10^{-11}$	294-300		
$1.95 \times 10^{-11} \exp[(125 \pm 20)/T]$	197-427	Dunlea and Ravishankara, 2004	PLP-RF (f)
$(3.00 \pm 0.24) \times 10^{-11}$	295		
$(3.29 \pm 0.27) \times 10^{-11}$	295	Takahashi et al., 2005	PLP-LIF (g)
$(3.3 \pm 0.6) \times 10^{-11}$	297-300	Dillon et al., 2008	PLP-LIF (h)

## Comments

- (a)  $\text{O}(^1\text{D})$  atoms were monitored by time-resolved detection of  $\text{O}(^1\text{D}) \rightarrow \text{O}(^3\text{P})$  emission.
- (b)  $\text{O}(^1\text{D})$  generated by  $\text{O}_3$  photolysis at 248 nm and  $\text{O}(^3\text{P})$  product monitored by RA. Pressures were 13- 26 mbar.
- (c)  $\text{O}(^1\text{D})$  generated by 266 nm photolysis of  $\text{O}_3$  and the  $\text{O}(^3\text{P})$  product monitored by RF. Most experiments conducted at 53 mbar (He bath gas).
- (d)  $\text{O}(^1\text{D})$  generated by the 193 nm photolysis of  $\text{N}_2\text{O}$  and monitored directly by VUV-LIF or indirectly, in a competitive kinetics method, as HO following reaction of  $\text{O}(^1\text{D})$  with  $\text{C}_4\text{H}_{10}$ . The competitive kinetics method yielded consistently large values of  $k$  (factor of 1.13). The value given in the table is from the VUV-LIF experiments only, as the authors suggest that this method is less prone to systematic error.
- (e)  $\text{O}(^1\text{D})$  generated by 248 or 266 nm photolysis of  $\text{O}_3$  and the  $\text{O}(^3\text{P})$  product monitored by RF. Most experiments conducted at  $\approx 27$  mbar (He bath gas).
- (f)  $\text{O}(^1\text{D})$  generated by 248 nm photolysis of  $\text{O}_3$  and the  $\text{O}(^3\text{P})$  product monitored by RF. Experiments conducted at 6.7 – 67 mbar (He bath gas).
- (g)  $\text{O}(^1\text{D})$  generated by the 193 nm photolysis of  $\text{N}_2\text{O}$  and monitored by VUV-LIF. Experiments at 6.7 mbar (He bath gas).

(h) Competitive kinetics method. O(<sup>1</sup>D) was generated in the 248 nm photolysis of O<sub>3</sub> and detected as HO following reaction with *n*-C<sub>6</sub>H<sub>14</sub> or H<sub>2</sub>O.

### Preferred Values

$k = 3.1 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at 298 K.

$k = 2.15 \times 10^{-11} \exp(110/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  over the temperature range 190-500 K.

### Reliability

$\Delta \log k = \pm 0.05$  at 298 K.


$\Delta(E/R) = \pm 30 \text{ K}$ .

### Comments on Preferred Values

Both the room temperature rate coefficient and values of *E/R* obtained for this reaction are in good agreement, although the datasets obtained prior to 2004 indicate slightly lower numbers for *k*. Three carefully executed, independent studies of this reaction in 2004 (Blitz et al., 2004; Strekowski and Wine, 2004; Dunlea and Ravishankara, 2004) using VUV-LIF detection of O(<sup>1</sup>D) and RF detection of O(<sup>3</sup>P) result in values of *k* that agree to within 5 % of an average value of  $3.1 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at room temperature and provide the basis of the preferred value at 298 K. The less comprehensive, room temperature studies of Takahashi et al (2005) and Dillon et al. (2008), which appeared subsequently, are consistent with this value. The preferred values are from Ravishankara et al. (2002) which combines the data from Blitz et al. (2004), Strekowski and Wine (2004) and Dunlea and Ravishankara (2004). Strekowski and Wine (2004) showed that the quenching rate coefficient for synthetic air is consistent with the individual values obtained for N<sub>2</sub> and O<sub>2</sub> with a ratio at 295 K of  $k(\text{O}(\text{}^1\text{D}) + \text{O}_2) / k(\text{O}(\text{}^1\text{D}) + \text{N}_2) = 1.30$ .

### References

- Amimoto, S. T., Force, A. P., Gulotty Jr. R. G., and Wiesenfeld, J. R.: J. Chem. Phys. 71, 3640, 1979.
- Blitz, M. A., Dillon, T. J., Heard, D. E., Pilling, M. J. and Trought, I. D.: Phys. Chem. Chem. Phys. 6, 2162-2171, 2004.
- Dillon, T. J., Horowitz, A., and Crowley, J. N.: Atmos. Chem. Phys. 8, 1547, 2008.
- Dunlea, E. J. and Ravishankara, A. R.: Phys. Chem. Chem. Phys. 6, 2152, 2004.
- Ravishankara, A. R., Dunlea, E. J., Blitz, M. A., Dillon, T. J., Heard, D. E., Pilling, M. J., Strekowski, R. S., Nicovich, J. M. and Wine, P. H.: Geophys. Res. Lett. 7, 861, 2002.
- Streit, G. E., Howard, C. J. Schmeltekopf, A. L., Davidson, J. A., and Schiff, H. I.: J. Chem. Phys. 65, 4761, 1976.
- Strekowski, R. S., Nicovich, J. M. and Wine, P. H.: Phys. Chem. Chem. Phys. 6, 2145, 2004.
- Takahashi, K., Takeuchi, Y. and Matsumi, Y.: Chem. Phys. Lett. 410, 196, 2005.
- Wine, P. H. and Ravishankara, A. R.: Chem. Phys. Lett. 77, 103, 1981.



**Data on  $k(\text{O}(^1\text{D}) + \text{N}_2)$  since 2004.**