

IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet HO_x_VOC79

Website: <http://iupac.pole-ether.fr>. See website for latest evaluated data. Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hard copy without explicit written permission. The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, <http://iupac.pole-ether.fr>.

This data sheet last evaluated: June 2011; last change in preferred values: June 2011.

HO₂ + CH₃CH(OH)CH(O₂)CH₃ → products

Rate coefficient data

<i>k</i> /cm ³ molecule ⁻¹ s ⁻¹	Temp./K	Reference	Technique/ Comments
<i>Absolute Rate Coefficients</i>			
(1.5 ± 0.4) × 10 ⁻¹¹	296	Jenkin and Hayman, 1995	PLP-UVA (a)

Comments

- (a) PLP-UV absorption study of H₂O₂-but-2-ene-O₂-N₂ mixtures at 1 bar (750 Torr) and 296 K. Conditions were chosen such that the initial [HO₂]/[CH₃CH(OH)CH(O₂)CH₃] was varied in the range 0.7 – 7.1. *k* was determined from simulation of transient decay traces recorded at 220 nm, 230 nm and 260 nm, with different relative contributions to the signals being due to CH₃CH(OH)CH(O₂)CH₃ and HO₂ absorption. No systematic dependence of the derived *k* on initial [HO₂]/[CH₃CH(OH)CH(O₂)CH₃] was observed.

Preferred Values

Parameter	Value	T/K
<i>k</i> /cm ³ molecule ⁻¹ s ⁻¹	1.5 × 10 ⁻¹¹	298
<i>Reliability</i>		
Δ log <i>k</i>	± 0.3	298

Comments on Preferred Values

The preferred rate coefficient is based on the sole kinetics study of Jenkin and Hayman (1995). Although the value of *k* is comparable with those reported for the reactions of HO₂ with other β-hydroxyalkyl peroxy radicals, HOCH₂CH₂O₂ (see datasheet HO_x_VOC55) and (CH₃)₂C(OH)C(O₂)(CH₃)₃ (Boyd et al., 2003), confirmatory studies are required to reduce the recommended uncertainty limits.

By analogy with the reaction of HO₂ with HOCH₂CH₂O₂, it is likely that the reaction mainly yields CH₃CH(OH)CH(OOH)CH₃ and O₂. Indeed, the API-MS study of Tuazon et al. (1998) reported formation of this hydroxyhydroperoxide, during the HO initiated oxidation of *cis*-but-2-ene and *trans*-but-2-ene in the absence of NO_x (at 297 K and 986 mbar), attributed to the reaction of CH₃CH(OH)CH(O₂)CH₃ with HO₂; although further product studies are required to quantify the branching ratio. Dillon and Crowley (2008) have investigated the

formation of HO from the reactions of HO₂ with selected RO₂ radicals. In the case of the structurally similar peroxy radicals, HOCH₂CH₂O₂ and CH₃CH(OH)CH₂O₂, respective upper limits of 0.04 and 0.06 were reported for the propagating channels forming HO, O₂ and the corresponding oxy radical. It is therefore probable that the title reaction is also mainly (or exclusively) terminating.

References

- Boyd, A. A., Flaud, P-M., Daugey, M. and Lesclaux, R.: J. Phys. Chem. A, 107, 818, 2003.
Dillon, T. J. and Crowley, J. N.: Atmos. Chem. Phys., 8, 4877, 2008.
Jenkin, M. E. and Hayman, G. D.: J. Chem. Soc. Faraday Trans., 91, 1911, 1995.
Tuazon, E.C., Aschmann, S.M., Arey, J. and Atkinson, R.: Environ. Sci. Technol., 32, 2106, 1998.