

# IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet NO<sub>3</sub>\_VOC34

Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hardcopy without explicit written permission. The citation for this data sheet is: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, (<http://iupac.pole-ether.fr>).

This data sheet last evaluated: June 2013; last change in preferred values: June 2013.



## Rate coefficient data

$k/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	Temp./K	Reference	Technique/ Comments
<i>Absolute Rate Coefficients</i>			
$(1.66 \pm 0.18) \times 10^{-11}$ $1.4 \times 10^{-12} \exp[(741 \pm 190)/T]$	298 298-433	Martínez et al., 1999	DF-LIF (a)
<i>Relative Rate Coefficients</i>			
$(1.87 \pm 0.11) \times 10^{-11}$ $(2.16 \pm 0.36) \times 10^{-11}$	295	Corchnoy and Atkinson, 1990	RR (b)

2-carene is 3,7,7-trimethyl-bicyclo[4.1.0]hept-2-ene.

## Comments

- (a) NO<sub>3</sub> radicals ( $6\text{--}30 \times 10^{11} \text{ molecule cm}^{-3}$ ) generated from reaction of F atoms (made in a microwave discharge through F<sub>2</sub>/He) with HNO<sub>3</sub>. Flow tube was operated at ~1.33 mbar (1 Torr) He at 4 temperatures between 298 and 433 K. 2-carene was present at similar concentrations (1-3 fold) to NO<sub>3</sub>. So that absolute NO<sub>3</sub> concentrations (derived by titration with tetramethylethane) were necessary to derive the rate coefficient.
- (b) 6400 L Teflon chamber at 295 K and 980 mbar (735 Torr) of air. NO<sub>3</sub> was generated by the thermal decomposition of N<sub>2</sub>O<sub>5</sub>. 2-carene and either 2,3-dimethyl-2-butene or 2-methyl-2-butene (reference reactants) were monitored by GC to obtain rate constant ratios,  $k(\text{NO}_3 + 2\text{-carene}) / k(\text{NO}_3 + 2,3\text{-dimethyl-2-butene}) = 0.377 \pm 0.063$  and  $k(\text{NO}_3 + 2\text{-carene}) / k(\text{NO}_3 + 2\text{-methyl-2-butene}) = 2.00 \pm 0.12$ . Using values of  $k(\text{NO}_3 + 2\text{-methyl-2-butene}) = 9.37 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at 298 K (Atkinson and Arey, 2003) and  $k(\text{NO}_3 + 2,3\text{-dimethyl-2-butene}) = 5.72 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  (Atkinson and Arey, 2003) the relative rates are converted to the absolute rate constants listed in the table. The value of  $(1.87 \pm 0.11) \times 10^{-11} \text{ molecule cm}^{-3}$  corresponds to the experiments in which 2-methyl-2-butene was used as reference.

## Preferred Values

Parameter	Value	T/K
-----------	-------	-----

$k / \text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	$2.0 \times 10^{-11}$	298
--	-----------------------	-----

*Reliability*

$\Delta \log k$	$\pm 0.12$	298
-----------------	------------	-----

*Comments on Preferred Values*

The three determinations of the room temperature rate coefficient agree to within ~30 % and the preferred value of the room temperature rate coefficient is based on the relative rate studies. The temperature dependence observed by Martínez et al. (1999) requires validation before a recommendation can be made.

There are no product studies of this reaction, though the large rate constant indicates that the reaction proceeds mainly via addition of  $\text{NO}_3$  across a double bond to form a chemically activated nitrooxyalkyl radical. At pressures found in the troposphere this adduct will undergo collisional stabilization prior to reaction with  $\text{O}_2$  to form a nitrooxyalkyl peroxy radical or decompose to release  $\text{NO}_2$ . At atmospheric pressure the formation of the peroxy radical will generally dominate.

## References

- Atkinson, R., and Arey, J., Chem. Rev., 103, 4605-4638, 2003.  
Corchnoy, S. B., and Atkinson, R., Env. Sci. Tech., 24, 1497-1502, 1990.  
Martínez, E., Cabañas, B., Aranda, A., Martín, P., and Salgado, S., J. Atmos. Chem., 33, 265-282, 1999.