

IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation – Data Sheet oRCIOx50

Website: <http://iupac.pole-ether.fr>. See website for latest evaluated data. Data sheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hardcopy without explicit written permission.

This data sheet updated: 27th January 2006.



Rate coefficient data ($k = k_1 + k_2$)

$k/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	Temp./K	Reference	Technique/ Comments
<i>Absolute Rate Coefficients</i>			
$k_{\text{obs}} = (4.13 \pm 0.58) \times 10^{-12}$	298	Wallington and Nielsen, 1991	FP-UVA (a,b)
<i>Branching Ratios</i>			
$k_2/k \approx 1.0$	298	Tuazon and Atkinson, 1994	UV-P-FTIR (e)

Comments

- (a) k_{obs} is based on the measured overall second-order decay of $\text{CF}_2\text{ClCH}_2\text{O}_2$, defined by $-\text{d}[\text{CF}_2\text{ClCH}_2\text{O}_2]/\text{dt} = 2k_{\text{obs}}[\text{CF}_2\text{ClCH}_2\text{O}_2]^2$. As described in detail by Lesclaux (1997), HO_2 radicals formed from the subsequent chemistry of $\text{CF}_2\text{ClCH}_2\text{O}$ (formed from channel (2)) are expected to lead to secondary removal of $\text{CF}_2\text{ClCH}_2\text{O}_2$. The true value of k is expected to fall in the range $k_{\text{obs}}/(1+\alpha) < k < k_{\text{obs}}$, where $\alpha = k_2/k$.
- (b) Pulse radiolysis study of $\text{CF}_2\text{ClCH}_3\text{-O}_2\text{-SF}_6$ mixtures over the pressure range 150-1000 mbar. $\text{CF}_2\text{ClCH}_2\text{O}_2$ radicals were monitored by UV absorption with $\sigma_{250 \text{ nm}} = (3.38 \pm 0.68) \times 10^{-18} \text{ cm}^2 \text{ molecule}^{-1}$.
- (c) Photolysis of Cl_2 in the presence of $\text{CF}_2\text{ClCH}_3\text{-air}$ mixtures at 987 mbar pressure. In situ monitoring of reactants and products by FTIR spectroscopy was consistent with formation of CF_2ClCHO with a yield *ca.* 100%, once corrections were made for secondary removal.

Preferred Values

$k = 2.8 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at 298 K.

$k_2/k = 1.0$ at 298 K.

Reliability

$\Delta \log k = \pm 0.4$ at 298 K.

$\Delta(k_2/k) = {}^{+0.0}_{-0.2}$ at 298 K.

Comments on Preferred Values

The observed formation of CF_2ClCHO in *ca.* 100% yield from CF_2ClCH_3 oxidation in the product study of Tuazon and Atkinson (1994) is consistent with the reaction proceeding predominantly *via* the radical forming channel (2), followed by reaction of $\text{CF}_2\text{ClCH}_2\text{O}$ with O_2 to form CF_2ClCHO and HO_2 . This also indicates that secondary removal of $\text{CF}_2\text{ClCH}_2\text{O}_2$ by reaction with HO_2 must generate CF_2ClCHO as the major carbon-containing product, in a similar fashion to the observed dominant formation of HC(O)Cl from the reaction of HO_2 with CH_2ClO_2 (Wallington et al., 1996).

The preferred value of k at 298 K is derived from the k_{obs} value reported by Wallington and

Nielsen (1991). Similar to a procedure adopted by Lesclaux (1997) for peroxy radicals for which the self reaction rate coefficients are \geq ca. $2 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, k is estimated to be $k_{\text{obs}}/(1+0.5(k_2/k))$, with this approximation assuming that the secondary reaction of HO_2 with $\text{CF}_2\text{ClCH}_2\text{O}_2$ competes equally with its removal via $\text{HO}_2 + \text{HO}_2$. The reliability range reflects that k has been derived by this approximate procedure. Confirmatory kinetics and product studies are required.

References

- Lesclaux, R.: Combination of peroxy radicals in the gas phase. In 'Peroxy Radicals', edited by Z.B. Alfassi. John Wiley and Sons, 1997.
- Tuazon, E.C. and Atkinson, R.: Environ. Sci. Technol., 28, 2306, 1994.
- Wallington, T. J. and Nielsen, O. J.: Int. J. Chem. Kinet., 23, 785, 1991.
- Wallington, T. J., Hurley, M. D. and Schneider, W. F.: Chem. Phys. Lett., 251, 164, 1996.